

Results	All Imaging (N=1,024)		Ultrasound (N=721)		CT Imaging (N=313)	
	Alone	+OVA1*	US Alone	+OVA1*	CT Alone	+OVA1*
Sensitivity	78.5% (153/195)	97.4% (190/195)	76.9% (70/91)	97.8% (89/91)	79.8% (83/104)	97.1% (101/104)
Specificity	53.8% (451/839)	28.5% (239/839)	55.4% (349/630)	30.8% (194/630)	48.8% (102/209)	21.5% (45/209)
Positive Predictive Value	28.3% (153/541)	24.1% (190/790)	19.9% (70/351)	17.0% (89/525)	43.7% (83/190)	38.1% (101/265)
Negative Predictive Value	91.5% (451/493)	98.0% (239/244)	94.3% (349/370)	99.0% (194/196)	82.9% (102/123)	93.8% (45/48)
Cancers Missed	21.5% (42/195)	2.6% (5/195)	23.3% (21/90)	2.2% (2/91)	20.2% (21/104)	2.9% (3/104)

Table 5. Results of study comparing sensitivity of imaging for detecting ovarian masses, alone and in combination with the OVA1 test. *Patient was classified as a high-risk result if imaging or OVA1 risk score were high risk, per OVA1 risk protocol.